

# Comment on “A note on the construction of the Ermakov-Lewis invariant”

F. Haas

Laboratório Nacional de Computação Científica  
Coordenação de Matemática Aplicada  
Av. Getúlio Vargas, 333  
25651-070, Petrópolis, RJ - Brazil  
ferhaas@lncc.br

J. Goedert

Centro de Ciências Exatas e Tecnológicas - UNISINOS  
Av. Unisinos, 950  
93022-000, São Leopoldo, RS - Brazil  
goedert@exatas.unisinos.br

## Abstract

We show that the basic results on the paper referred in the title [J. Phys. A: Math. Gen. **35** (2002) 5333-5345], concerning the derivation of the Ermakov invariant from Noether symmetry methods, are not new.

*PACS numbers: 02.30.Hg, 02.90.+p, 03.20.+i*

The purpose of this comment is to point out that the main results presented in a recently published paper [1], are not new. At the end of the introduction of this paper, the authors claims that “... *this is the first time the Noether symmetries are being considered to discuss the source of the Ermakov-Lewis*

*invariant.*” Unfortunately, the authors miss the reference *Dynamical symmetries and the Ermakov invariant*, by Haas and Goedert [2]. In this paper, the Ermakov invariant is, apparently for the first time, deduced as a consequence of a dynamical Noether symmetry. To make this point clear, it is enough to compare equation (29) in the paper by Haas and Goedert with Proposition 1 and equation (4.17) in the paper by Moyo and Leach. Both equations present the dynamical symmetry associated to the Ermakov invariant, for the case of Lagrangian Ermakov systems, as the result from a straightforward application of the converse of Noether’s theorem.

We further notice that the Lagrangian formulation for Ermakov systems in the referred publication [1] is by no means new. This can be seen by comparing the potential functions from equations (11) in the work by Haas and Goedert with the potential given by equation (3.18) in the work by Moyo and Leach. This later work also ignores the Hamiltonian descriptions for Ermakov systems developed earlier in [3], for the case of frequency functions depending on time only, and in [4], for the case of frequency functions depending also on dynamical variables.

As a final remark, the work by Moyo and Leach also ignores the papers [5]-[8], dedicated to the analysis of uncoupled Ermakov systems in the light of Noether’s theorem. These works, however, do not deal with truly two-dimensional, coupled Ermakov systems, as in the case of references [1, 2]. Rather, these papers [5]-[8] deal with Ermakov systems in which one of the equations plays the principal role, while the other, decoupled from the first, is treated as an auxiliary equation. In these cases, the Lagrangian description is effectively one-dimensional and the Ermakov invariant cannot be obtained as a result from an associated dynamical Noether symmetry.

### **Acknowledgements**

This work has been supported by the Brazilian agency Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

### **References**

- [1] Moyo S and Leach P G L 2002 A note on the construction of the Ermakov-Lewis invariant *J. Phys. A: Math. Gen.* **35** 5333-5345

- [2] Haas F and Goedert J 2001 Dynamical symmetries and the Ermakov invariant *Phys. Lett. A* **279** 181-188
- [3] Cerveró J M and Lejarreta J D 1991 Ermakov Hamiltonians *Phys. Lett. A* **156** 201-205
- [4] Haas F and Goedert J 1996 On the Hamiltonian structure of Ermakov systems *J. Phys. A: Math. Gen.* **29** 4083-4092
- [5] Ray J R and Reid J L 1979 Noether's theorem, time-dependent invariants and nonlinear equations of motion *J. Math. Phys.* **20** 2054-2057
- [6] Ray J R 1981 Invariants for nonlinear equations of motion *Prog. Theor. Phys.* **65** 877-882
- [7] Ray J R 1980 Noether's theorem and exact invariants for time-dependent systems *J. Phys. A: Math. Gen* **13** 1969-1975
- [8] Kaushal R S and Korsch H J 1981 Dynamical Noether invariants for time-dependent nonlinear systems *J. Math. Phys.* **22** 1904-1908