

Thus  $c_1 = 2, c_2 = 3$ , and  $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ . The coordinate system on  $H$  determined by  $\mathcal{B}$  is shown in Figure 7. ■




FIGURE 7 A coordinate system on a plane  $H$  in  $\mathbb{R}^3$ .

If a different basis for  $H$  were chosen, would the associated coordinate system also make  $H$  isomorphic to  $\mathbb{R}^2$ ? Surely, this must be true. We shall prove it in the next section.

**PRACTICE PROBLEMS**

1. Let  $\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} -3 \\ 4 \\ 0 \end{bmatrix}, \mathbf{b}_3 = \begin{bmatrix} 3 \\ -6 \\ 3 \end{bmatrix}$ , and  $\mathbf{x} = \begin{bmatrix} -8 \\ 2 \\ 3 \end{bmatrix}$ .
  - a. Show that the set  $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$  is a basis of  $\mathbb{R}^3$ .
  - b. Find the change-of-coordinates matrix from  $\mathcal{B}$  to the standard basis.
  - c. Write the equation that relates  $\mathbf{x}$  in  $\mathbb{R}^3$  to  $[\mathbf{x}]_{\mathcal{B}}$ .
  - d. Find  $[\mathbf{x}]_{\mathcal{B}}$ , for the  $\mathbf{x}$  given above.
2. The set  $\mathcal{B} = \{1 + t, 1 + t^2, t + t^2\}$  is a basis for  $\mathbb{P}_2$ . Find the coordinate vector of  $\mathbf{p}(t) = 6 + 3t - t^2$  relative to  $\mathcal{B}$ .

**4.4 EXERCISES**

In Exercises 1–4, find the vector  $\mathbf{x}$  determined by the given coordinate vector  $[\mathbf{x}]_{\mathcal{B}}$  and the given basis  $\mathcal{B}$ .

1.  $\mathcal{B} = \left\{ \begin{bmatrix} 3 \\ -5 \end{bmatrix}, \begin{bmatrix} -4 \\ 6 \end{bmatrix} \right\}, [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 5 \\ 3 \end{bmatrix}$
2.  $\mathcal{B} = \left\{ \begin{bmatrix} 4 \\ 5 \end{bmatrix}, \begin{bmatrix} 6 \\ 7 \end{bmatrix} \right\}, [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 8 \\ -5 \end{bmatrix}$
3.  $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -4 \\ 3 \end{bmatrix}, \begin{bmatrix} 5 \\ 2 \\ -2 \end{bmatrix}, \begin{bmatrix} 4 \\ -7 \\ 0 \end{bmatrix} \right\}, [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix}$
4.  $\mathcal{B} = \left\{ \begin{bmatrix} -1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ -5 \\ 2 \end{bmatrix}, \begin{bmatrix} 4 \\ -7 \\ 3 \end{bmatrix} \right\}, [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} -4 \\ 8 \\ -7 \end{bmatrix}$

In Exercises 5–8, find the coordinate vector  $[\mathbf{x}]_{\mathcal{B}}$  of  $\mathbf{x}$  relative to the given basis  $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ .

5.  $\mathbf{b}_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 2 \\ -5 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$
6.  $\mathbf{b}_1 = \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 5 \\ -6 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 4 \\ 0 \end{bmatrix}$
7.  $\mathbf{b}_1 = \begin{bmatrix} 1 \\ -1 \\ -3 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} -3 \\ 4 \\ 9 \end{bmatrix}, \mathbf{b}_3 = \begin{bmatrix} 2 \\ -2 \\ 4 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 8 \\ -9 \\ 6 \end{bmatrix}$
8.  $\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \\ 8 \end{bmatrix}, \mathbf{b}_3 = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 3 \\ -5 \\ 4 \end{bmatrix}$

In Exercises 9 and 10, find the change-of-coordinates matrix from  $\mathcal{B}$  to the standard basis in  $\mathbb{R}^n$ .

9.  $\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ -9 \end{bmatrix}, \begin{bmatrix} 1 \\ 8 \end{bmatrix} \right\}$
10.  $\mathcal{B} = \left\{ \begin{bmatrix} 3 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -5 \end{bmatrix}, \begin{bmatrix} 8 \\ -2 \\ 7 \end{bmatrix} \right\}$

In Exercises 11 and 12, use an inverse matrix to find  $[\mathbf{x}]_{\mathcal{B}}$  for the given  $\mathbf{x}$  and  $\mathcal{B}$ .

11.  $\mathcal{B} = \left\{ \begin{bmatrix} 3 \\ -5 \end{bmatrix}, \begin{bmatrix} -4 \\ 6 \end{bmatrix} \right\}, \mathbf{x} = \begin{bmatrix} 2 \\ -6 \end{bmatrix}$
12.  $\mathcal{B} = \left\{ \begin{bmatrix} 4 \\ 5 \end{bmatrix}, \begin{bmatrix} 6 \\ 7 \end{bmatrix} \right\}, \mathbf{x} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$

13. The set  $\mathcal{B} = \{1 + t^2, t + t^2, 1 + 2t + t^2\}$  is a basis for  $\mathbb{P}_2$ . Find the coordinate vector of  $\mathbf{p}(t) = 1 + 4t + 7t^2$  relative to  $\mathcal{B}$ .
14. The set  $\mathcal{B} = \{1 - t^2, t - t^2, 2 - 2t + t^2\}$  is a basis for  $\mathbb{P}_2$ . Find the coordinate vector of  $\mathbf{p}(t) = 3 + t - 6t^2$  relative to  $\mathcal{B}$ .

In Exercises 15 and 16, mark each statement True or False. Justify each answer. Unless stated otherwise,  $\mathcal{B}$  is a basis for a vector space  $V$ .

15. a. If  $\mathbf{x}$  is in  $V$  and if  $\mathcal{B}$  contains  $n$  vectors, then the  $\mathcal{B}$ -coordinate vector of  $\mathbf{x}$  is in  $\mathbb{R}^n$ .  
 b. If  $P_{\mathcal{B}}$  is the change-of-coordinates matrix, then  $[\mathbf{x}]_{\mathcal{B}} = P_{\mathcal{B}}\mathbf{x}$ , for  $\mathbf{x}$  in  $V$ .  
 c. The vector spaces  $\mathbb{P}_3$  and  $\mathbb{R}^3$  are isomorphic.
16. a. If  $\mathcal{B}$  is the standard basis for  $\mathbb{R}^n$ , then the  $\mathcal{B}$ -coordinate vector of an  $\mathbf{x}$  in  $\mathbb{R}^n$  is  $\mathbf{x}$  itself.  
 b. The correspondence  $[\mathbf{x}]_{\mathcal{B}} \mapsto \mathbf{x}$  is called the coordinate mapping.  
 c. In some cases, a plane in  $\mathbb{R}^3$  can be isomorphic to  $\mathbb{R}^2$ .
17. The vectors  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 2 \\ -8 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -3 \\ 7 \end{bmatrix}$  span  $\mathbb{R}^2$  but do not form a basis. Find two different ways to express  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  as a linear combination of  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ .
18. Let  $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$  be a basis for a vector space  $V$ . Explain why the  $\mathcal{B}$ -coordinate vectors of  $\mathbf{b}_1, \dots, \mathbf{b}_n$  are the columns  $\mathbf{e}_1, \dots, \mathbf{e}_n$  of the  $n \times n$  identity matrix.
19. Let  $S$  be a finite set in a vector space  $V$  with the property that every  $\mathbf{x}$  in  $V$  has a unique representation as a linear combination of elements of  $S$ . Show that  $S$  is a basis of  $V$ .
20. Suppose  $\{\mathbf{v}_1, \dots, \mathbf{v}_4\}$  is a linearly dependent spanning set for a vector space  $V$ . Show that each  $\mathbf{w}$  in  $V$  can be expressed in more than one way as a linear combination of  $\mathbf{v}_1, \dots, \mathbf{v}_4$ . [Hint: Let  $\mathbf{w} = k_1\mathbf{v}_1 + \dots + k_4\mathbf{v}_4$  be an arbitrary vector in  $V$ .

Use the linear dependence of  $\{\mathbf{v}_1, \dots, \mathbf{v}_4\}$  to produce another representation of  $\mathbf{w}$  as a linear combination of  $\mathbf{v}_1, \dots, \mathbf{v}_4$ .]

21. Let  $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -4 \end{bmatrix}, \begin{bmatrix} -2 \\ 9 \end{bmatrix} \right\}$ . Since the coordinate mapping determined by  $\mathcal{B}$  is a linear transformation from  $\mathbb{R}^2$  into  $\mathbb{R}^2$ , this mapping must be implemented by some  $2 \times 2$  matrix  $A$ . Find it. [Hint: Multiplication by  $A$  should transform a vector  $\mathbf{x}$  into its coordinate vector  $[\mathbf{x}]_{\mathcal{B}}$ .]
22. Let  $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$  be a basis for  $\mathbb{R}^n$ . Produce a description of an  $n \times n$  matrix  $A$  that implements the coordinate mapping  $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ . (See Exercise 21.)

Exercises 23–26 concern a vector space  $V$ , a basis  $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ , and the coordinate mapping  $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ .

23. Show that the coordinate mapping is one-to-one. [Hint: Suppose  $[\mathbf{u}]_{\mathcal{B}} = [\mathbf{w}]_{\mathcal{B}}$  for some  $\mathbf{u}$  and  $\mathbf{w}$  in  $V$ , and show that  $\mathbf{u} = \mathbf{w}$ .]
24. Show that the coordinate mapping is onto  $\mathbb{R}^n$ . That is, given any  $\mathbf{y}$  in  $\mathbb{R}^n$ , with entries  $y_1, \dots, y_n$ , produce  $\mathbf{u}$  in  $V$  such that  $[\mathbf{u}]_{\mathcal{B}} = \mathbf{y}$ .
25. Show that a subset  $\{\mathbf{u}_1, \dots, \mathbf{u}_p\}$  in  $V$  is linearly independent if and only if the set of coordinate vectors  $\{[\mathbf{u}_1]_{\mathcal{B}}, \dots, [\mathbf{u}_p]_{\mathcal{B}}\}$  is linearly independent in  $\mathbb{R}^n$ . [Hint: Since the coordinate mapping is one-to-one, the following equations have the same solutions,  $c_1, \dots, c_p$ .]

$$c_1\mathbf{u}_1 + \dots + c_p\mathbf{u}_p = \mathbf{0} \quad \text{The zero vector in } V$$

$$[c_1\mathbf{u}_1 + \dots + c_p\mathbf{u}_p]_{\mathcal{B}} = [\mathbf{0}]_{\mathcal{B}} \quad \text{The zero vector in } \mathbb{R}^n$$

26. Given vectors  $\mathbf{u}_1, \dots, \mathbf{u}_p$ , and  $\mathbf{w}$  in  $V$ , show that  $\mathbf{w}$  is a linear combination of  $\mathbf{u}_1, \dots, \mathbf{u}_p$  if and only if  $[\mathbf{w}]_{\mathcal{B}}$  is a linear combination of the coordinate vectors  $[\mathbf{u}_1]_{\mathcal{B}}, \dots, [\mathbf{u}_p]_{\mathcal{B}}$ .

In Exercises 27–30, use coordinate vectors to test the linear independence of the sets of polynomials. Explain your work.

27.  $1 + 2t^3, 2 + t - 3t^2, -t + 2t^2 - t^3$
28.  $1 - 2t^2 - t^3, t + 2t^3, 1 + t - 2t^2$
29.  $(1 - t)^2, t - 2t^2 + t^3, (1 - t)^3$
30.  $(2 - t)^3, (3 - t)^2, 1 + 6t - 5t^2 + t^3$
31. Use coordinate vectors to test whether the following sets of polynomials span  $\mathbb{P}_2$ . Justify your conclusions.  
 a.  $1 - 3t + 5t^2, -3 + 5t - 7t^2, -4 + 5t - 6t^2, 1 - t^2$   
 b.  $5t + t^2, 1 - 8t - 2t^2, -3 + 4t + 2t^2, 2 - 3t$
32. Let  $\mathbf{p}_1(t) = 1 + t^2, \mathbf{p}_2(t) = t - 3t^2, \mathbf{p}_3(t) = 1 + t - 3t^2$ .  
 a. Use coordinate vectors to show that these polynomials form a basis for  $\mathbb{P}_2$ .  
 b. Consider the basis  $\mathcal{B} = \{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$  for  $\mathbb{P}_2$ . Find  $\mathbf{q}$  in  $\mathbb{P}_2$ , given that  $[\mathbf{q}]_{\mathcal{B}} = \begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix}$ .

## 226 CHAPTER 4 Vector Spaces

In Exercises 33 and 34, determine whether the sets of polynomials form a basis for  $\mathbb{P}_3$ . Justify your conclusions.

33. [M]  $3 + 7t, 5 + t - 2t^3, t - 2t^2, 1 + 16t - 6t^2 + 2t^3$

34. [M]  $5 - 3t + 4t^2 + 2t^3, 9 + t + 8t^2 - 6t^3, 6 - 2t + 5t^2, t^3$

35. [M] Let  $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$  and  $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2\}$ . Show that  $\mathbf{x}$  is in  $H$  and find the  $\mathcal{B}$ -coordinate vector of  $\mathbf{x}$ , for

$$\mathbf{v}_1 = \begin{bmatrix} 11 \\ -5 \\ 10 \\ 7 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 14 \\ -8 \\ 13 \\ 10 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 19 \\ -13 \\ 18 \\ 15 \end{bmatrix}$$


36. [M] Let  $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  and  $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ . Show that  $\mathcal{B}$  is a basis for  $H$  and  $\mathbf{x}$  is in  $H$ , and find the  $\mathcal{B}$ -coordinate vector of  $\mathbf{x}$ , for

$$\mathbf{v}_1 = \begin{bmatrix} -6 \\ 4 \\ -9 \\ 4 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 8 \\ -3 \\ 7 \\ -3 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -9 \\ 5 \\ -8 \\ 3 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 4 \\ 7 \\ -8 \\ 3 \end{bmatrix}$$

[M] Exercises 37 and 38 concern the crystal lattice for titanium, which has the hexagonal structure shown on the left in the accompanying figure. The vectors

forming a basis for the unit cell shown on the right. The numbers here are Ångstrom units ( $1 \text{ \AA} = 10^{-8} \text{ cm}$ ). In alloys of titanium,

some additional atoms may be in the unit cell at the *octahedral* and *tetrahedral* sites (so named because of the geometric objects formed by atoms at these locations).



The hexagonal close-packed lattice and its unit cell.

37. One of the octahedral sites is  $\begin{bmatrix} 1/2 \\ 1/4 \\ 1/6 \end{bmatrix}$ , relative to the lattice basis. Determine the coordinates of this site relative to the standard basis of  $\mathbb{R}^3$ .

38. One of the tetrahedral sites is  $\begin{bmatrix} 1/2 \\ 1/2 \\ 1/3 \end{bmatrix}$ . Determine the coordinates of this site relative to the standard basis of  $\mathbb{R}^3$ .

## SOLUTIONS TO PRACTICE PROBLEMS

1. a. It is evident that the matrix  $P_{\mathcal{B}} = [\mathbf{b}_1 \ \mathbf{b}_2 \ \mathbf{b}_3]$  is row-equivalent to the identity matrix. By the Invertible Matrix Theorem,  $P_{\mathcal{B}}$  is invertible and its columns form a basis for  $\mathbb{R}^3$ .

b. From part (a), the change-of-coordinates matrix is  $P_{\mathcal{B}} = \begin{bmatrix} 1 & -3 & 3 \\ 0 & 4 & -6 \\ 0 & 0 & 3 \end{bmatrix}$ .

c.  $\mathbf{x} = P_{\mathcal{B}}[\mathbf{x}]_{\mathcal{B}}$

d. To solve the equation in (c), it is probably easier to row reduce an augmented matrix than to compute  $P_{\mathcal{B}}^{-1}$ :

$$\begin{bmatrix} 1 & -3 & 3 & -8 \\ 0 & 4 & -6 & 2 \\ 0 & 0 & 3 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & -5 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 1 \end{bmatrix}$$

$P_{\mathcal{B}} \quad \mathbf{x} \quad I \quad [\mathbf{x}]_{\mathcal{B}}$

Hence

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} -5 \\ 2 \\ 1 \end{bmatrix}$$

2. The coordinates of  $\mathbf{p}(t) = 6 + 3t - t^2$  with respect to  $\mathcal{B}$  satisfy

$$c_1(1+t) + c_2(1+t^2) + c_3(t+t^2) = 6 + 3t - t^2$$